

$\sin^2 \theta_W$ theory and new physics

LEE Hye-Sung

(Center for Theoretical Physics of the Universe, IBS, Daejeon 34051, Korea)

Abstract: After briefly discussing the importance of the precise measurement of the weak mixing angle, the implication of the dark Z on the low- Q^2 parity tests is discussed. The dark Z is a very light (roughly, MeV-GeV scale) gauge boson, which couples to the electromagnetic current as well as the weak neutral current.

Key words: parity test; dark force; dark photon; dark Z

CLC number: O572.3 **Document code:** A doi:10.3969/j.issn.0253-2778.2016.06.004

Citation: LEE H S. $\sin^2 \theta_W$ theory and new physics[J]. Journal of University of Science and Technology of China, 2016, 46(6): 470-475.

LEE H S. $\sin^2 \theta_W$ 理论与新物理[J]. 中国科学技术大学学报, 2016, 46(6): 470-475.

$\sin^2 \theta_W$ 理论与新物理

LEE Hye-Sung

(理论物理中心, 大田市 34051, 韩国)

摘要: 简要介绍弱混合角的精确测量的重要意义后, 讨论了在低能区宇称检验中的暗 Z 粒子的意义. 暗 Z 粒子是一种非常轻(大致在 MeV 到 GeV 的尺度)的规范玻色子, 能够在耦合电磁流的同时耦合弱中性流.

关键词: 宇称检验; 暗作用力; 暗光子; 暗 Z 粒子

0 Introduction

In this article, we emphasize the importance of the low- Q^2 parity test for the new physics searches. We illustrate our point with a specific example called the dark parity violation^[1-4], which means the parity violation induced by a dark gauge boson. This presentation shares some parts with Ref. [5], although updates and complementary descriptions are provided.

Let us briefly look back on the history of the $\sin^2 \theta_W$ physics. It is well documented in the review^[6], and we will go over only some part of it very briefly. In

1961, Glashow introduced the $SU(2)_L \times U(1)_L$ symmetry, which has a mixing between two neutral gauge bosons^[7]. In 1967, Weinberg added the Higgs mechanism with a Higgs doublet and a vacuum expectation value, establishing the mass relation $m_W = m_Z \cos \theta_W$ with the weak mixing angle θ_W ^[8]. He also predicted the weak neutral current mediated by the Z boson. In 1973, the neutral current was discovered in the neutrino scattering experiments at the CERN Gargamelle detector^[9]. Whether the $SU(2)_L \times U(1)_Y$ is a correct theory to describe this neutral current was not clear then though. One of the features of the

Received: 2015-11-30; **Revised:** 2016-04-20

Foundation item: Supported in part by the IBS(IBS-R018-D1).

Biography: LEE Hye-Sung(corresponding author), Professor/PhD. Research field: high energy physics. E-mail: hlee@ibs.re.kr

$SU(2)_L \times U(1)_Y$ was the mixing term in the weak neutral current interaction, proportional to $\sin^2\theta_w$, and the parity test measuring this $\sin^2\theta_w$ can possibly test the standard model (SM).

In 1978, SLAC E122 experiment using the polarized electron beam and the deuteron target measured the parity violation asymmetry, which gave $\sin^2\theta_w \approx 0.22$ (2), agreeing to the SM^[10]. It is noticeable that this establishment of the $SU(2)_L \times U(1)_Y$ by the SLAC parity test in 1978 occurred much earlier than the direct discovery of the W/Z boson resonances at the CERN SPS experiments in 1983^[11-12]. In 1979, after only one year of the SLAC parity test, Glashow, Salam, and Weinberg received the Nobel prize in physics.

The lessons we can learn from this history include: ① the parity test (by the precise measurement of $\sin^2\theta_w$) can be a critical way to search for a new gauge interaction; ② its finding may precede the direct discovery of a gauge boson by the bump search.

Fig. 1 taken from Ref. [4] shows the running of the $\sin^2\theta_w$ in the SM and the current experimental constraints. While the current data are more or less consistent with the SM prediction with the given error bars, more precise measurements in the future experiments (red bars) may reveal potential new physics effects that were elusive for the current constraints.




Fig. 1 The running of the $\sin^2\theta_w$

In Fig. 1, the running of the $\sin^2\theta_w$ with the momentum transfer Q in the SM and the current experimental constraints taken from Ref. [4]. The red bars show the anticipated sensitivities in the future

parity tests.

1 Dark Photon vs. Dark Z

The dark gauge boson (we use Z' for its notation) is a hypothetical particle with a very small mass and a small coupling to the SM particles. While the heavy Z' (typically TeV scale) has been a traditional target of discovery^[13], the light Z' (typically MeV-GeV scale) is a recently highlighted subject with a growing interest^[14]. For such a light particle to survive all experimental constraints, it should have extremely small couplings to the SM particles.

There are number of dark force models in the literature, but we consider only two of them. Both models commonly assume the kinetic mixing between the $U(1)_Y$ and the $U(1)_{\text{dark}}$ ^[15]. The SM particles have zero charges under the new gauge group $U(1)_{\text{dark}}$, yet the gauge boson Z' of the $U(1)_{\text{dark}}$ can still couple to the SM fermions through the mixing with the SM gauge bosons.

One model is the dark photon^[16], which couples only to the electromagnetic current at the leading order. Another is a relatively new model, the dark Z^[1], which couples to the electromagnetic current as well as the weak neutral current. Their interactions are given by

$$\mathcal{L}_{\text{dark } \gamma} = -\epsilon e J_{EM}^\mu Z'_{\mu} \quad (1)$$

$$\mathcal{L}_{\text{dark } Z} = -[\epsilon e J_{EM}^\mu + \epsilon_Z (g/2 \cos\theta_w) J_{NC}^\mu] Z'_{\mu} \quad (2)$$

with $J_{\mu}^{\text{EM}} = Q_f \bar{f} \gamma_{\mu} f$ and $J_{\mu}^{\text{NC}} = (T_{3f} - 2Q_f \sin^2\theta_w) \bar{f} \cdot \gamma_{\mu} f - (T_{3f}) \bar{f} \gamma_{\mu} \gamma_5 f$. ϵ and ϵ_Z are the parametrization of the effective γ - Z' mixing and Z - Z' mixing, respectively.

The difference of the two models comes from how the Z' gets a mass or the details of the Higgs sector. Because of the Z coupling, the Z' in the dark Z model inherits some properties of the Z boson such as the parity violating nature. In a rough sense, the dark photon is a heavier version of the photon, and the dark Z is a lighter version of the Z boson.

Because of the new coupling, some experiments that are irrelevant to the dark photon searches are relevant to the dark Z searches^[1-4, 17-20]. They include

the low - Q^2 parity test, which will be discussed later in this article.

2 Bump hunt

There are many ongoing and proposed searches for the dark force in the labs around the world^[14]. A particularly attractive feature about the dark force is that it is one of the rare new physics scenarios that can be tested/discovered at the low-energy experiments, which are typically built for nuclear or hadronic physics. Of course, it is possible because the dark force carrier Z' is very light (MeV-GeV scale).

Fig. 2 in Ref. [21] shows the parameter space of the dark photon with the current bounds. The bounds come from the electron^[22-23] and muon^[24-26] anomalous magnetic moments, fixed target experiments^[27-28], beam dump experiments^[29], meson decays^[30-35], and e^+e^- collision ($e^+e^- \rightarrow \gamma + l^+l^-$) experiments^[36-37]. Except for the anomalous magnetic moments, the searches are all based on the dilepton searches from the Z' , that is the bump hunt.




Fig. 2 Dark force searches all around the world

If we put some of these experimental efforts on the map (Fig. 2), we can see the search is practically a global activity. Quite obviously, we are going through a very exciting time with so many contemporary searches to find a new fundamental force of nature.

3 Low-energy parity test

Now, we discuss the low - energy parity test^[1-4] as another means to search for the dark force. The presence of the dark Z modifies the effective lagrangian of the weak neutral current scattering,

$$\mathcal{L}_{\text{eff}} = -\frac{4G_F}{\sqrt{2}} J_{\text{NC}}^\mu(\sin^2\theta_W) J_\mu^{\text{NC}}(\sin^2\theta_W) \quad (3)$$

$$G_F \rightarrow \left(1 + \delta^2 \frac{1}{1 + Q^2/m_{Z'}^2}\right) G_F \quad (4)$$

$$\sin^2\theta_W \rightarrow \left(1 - \varepsilon\delta \frac{m_Z}{m_{Z'}} \frac{\cos\theta_W}{\sin\theta_W} \frac{1}{1 + Q^2/m_{Z'}^2}\right) \sin^2\theta_W \quad (5)$$

where Q is the momentum transfer between the two neutral currents, and δ is a reparametrization of the ε_Z with $\varepsilon_Z = (m_{Z'}/m_Z)\delta$. One salient feature is that these shifts are sensitive only to the low- Q^2 (low momentum transfer). Thus, the dark Z effectively changes the weak neutral current scattering, including the effective $\sin^2\theta_W$, which describes the parity violation, but only for the low momentum transfer.

Fig. 3 (from Ref. [3] with a slight modification) shows an example of how the effective $\sin^2\theta_W$ changes with Q in the presence of a dark Z for $m_{Z'} = 100$ MeV, 200 MeV cases. Although there are some details in the figure, the important point is that the deviations appear only in the low Q values, roughly $Q \leq m_{Z'}$. They never appear in the high Q values relevant to the high-energy experiments, which tells us that we need low-energy experiments to see the dark Z mediated scattering effects.




Fig. 3 Effective $\sin^2\theta_W$ running taken from Ref. [3]

In Fig. 3, dark Z of 100 MeV and 200 MeV were taken. Note that the deviations appear only in the low- Q^2 regime ($Q^2 \leq m_{Z'}^2$).

In this region, non-perturbative QCD contributions to the SM value become important. They have traditionally been determined utilizing dispersion relations^[38-39]. Recently, first-principle lattice QCD

determinations of the leading-order hadronic effects have also become available^[40-41].

For the low- Q^2 parity tests, one can use the atomic parity violation in Cs^[42-44], Ra⁺ ion^[45-46] or the low- Q^2 polarized electron scattering experiments SLAC E158^[47], JLAB Q_{weak} ^[48], JLAB Møller^[49] and Mainz P2^[50]. The possible deviations due to the dark Z can be large enough to be observed with the future experiments.

For the intermediate scale Z' of $m_{Z'} \approx \mathcal{O}(10)$ Gev, the deep inelastic scattering experiments such as JLAB PVDIS^[51] and JLAB SOLID^[52] may also be sensitive. In fact, as Fig. 4 (The NuTeV anomaly can be addressed by this intermediate scale dark Z) taken from Ref. [4] shows, the intermediate scale Z' can address the NuTeV ($\langle Q \rangle \approx 5$ Gev) anomaly^[53].




Fig. 4 The 15 GeV dark Z case taken from Ref. [4]

4 Conclusion

The parity test by precise measurement of the $\sin^2\theta_w$ has been important in studying new gauge interactions. Especially, it critically helped establishing the $SU(2)_L \times U(1)_Y$ electroweak theory. There is a growing interest in the dark gauge interaction (mediated by a light Z' gauge boson) around the world partly because many existing low-energy facilities can join the searches. While most searches of the light Z' are based on the direct bump searches, the parity tests in the low- Q^2 (such as the atomic parity violation, polarized electron scattering, deep inelastic scattering) are important and complementary searches for the dark force. The latter are also independent of the Z' decay branching ratios.

If the history may repeat, the dark force evidence

from the low- Q^2 parity test might precede the discovery of a new resonance just like what happened in the electroweak interaction case.

References

- [1] DAVOUDIASL, LEE H S, MARCIANO W J. "Dark" Z implications for parity violation, rare meson decays, and Higgs physics[J]. Physical Review D, 2012, 85(11): 115019(1-13).
- [2] DAVOUDIASLH, LEE H S, MARCIANO W J. Muon anomaly and dark parity violation[J]. Physical Review Letters, 2012, 109(3): 031802(1-5).
- [3] DAVOUDIASLH, LEE H S, MARCIANO W J. Muon $g-2$, rare kaon decays, and parity violation from dark bosons[J]. Physical Review D, 2014, 89(9): 98-105.
- [4] DAVOUDIASL H, LEE H S, MARCIANO W J. Low Q^2 weak mixing angle measurements and rare higgs decays [J]. Physical Review D, 2015, 92(5): 055005(1-5).
- [5] KLOPOTH S. Parity violation by a dark gauge boson[J]. Eprint Arxiv, 2014: arXiv:1410.8435.
- [6] KUMARK S, MANTRY S, MARCIANO W J, et al. Low energy measurements of the weak mixing angle [J]. Annual Review of Nuclear and Particle Science, 2013, 63(6): 237-267.
- [7] GLASHOW S L. Partial-symmetries of weak interactions [J]. Nuclear Physics, 1961, 22(4): 579-588.
- [8] WEINBERG S. A model of leptons[J]. Physical Review Letters, 1967, 19: 1264-1266.
- [9] HASERT F J, KABE S, KRENZ W, et al. Observation of neutrino-like interactions without muon or electron in the Gargamelle neutrino experiment [J]. Physcs Letters B, 1973, 46(1): 138-140.
- [10] PRESCOTT C Y, ATWOOD W B, COTTRELL R L A, et al. Parity non-conservation in inelastic electron scattering [J]. Physics Letters B, 1978, 77(3): 347-352.
- [11] BANNER M, BATTISTON R, BLOCH P, et al. Observation of single isolated electrons of high transverse momentum in events with missing transverse energy at the CERN $\bar{p}p$ collider[J]. Physics Letters B, 1983, 122(5-6): 476-485.
- [12] ARNISON G, ASTBURY A, AUBERT B, et al. Experimental-observation of lepton pairs of invariant mass around 95 GeV/ c^2 at the cern sps collider[J]. Physics Letters B, 1983, 126: 398-410.
- [13] LANGACKER P. The physics of heavy Z' gauge bosons

- [J]. Review of Modern Physics, 2009, 81 (3): 1199-1228.
- [14] ESSIG R, JAROS J A, WESTER W, et al. Dark sectors and new, light, weakly-coupled particles [J]. Eprint Arxiv, 2013; arXiv:1311.0029.
- [15] HOLDOM B. Two $U(1)$'s and ε charge shifts [J]. Physics Letters B, 1986, 166(2): 196-198.
- [16] ARKANI-HAMED N, FINKBEINER D P, SLATYER T R, ET AL. A theory of dark matter [J]. Physical Review D, 2009, 79(1): 015014(1-22).
- [17] DAVOUDIASHVILI, LEE H S, LEWIS I et al. Higgs decays as a window into the dark sector [J]. Physical Review D, 2013, 88(1): 392-426.
- [18] DAVOUDIASHVILI H, MARCIANO W J, RAMOS R, et al. Charged higgs discovery in the W plus "Dark" vector boson decay mode [J]. Physics, 2014, 89 (1): 1858-1871.
- [19] KONG K, LEE H S, PARK M. Dark decay of the top quark [J]. Physical Review D, 2014, 89(7): 539-560.
- [20] KIM D, LEE H S, PARK M. Invisible dark gauge boson search in top decays using a kinematic method [J]. Journal of High Energy Physics, 2015, 134 (3): 1-2 (arXiv:1411.0668).
- [21] SOFFER A. Searches for light scalars, pseudoscalars, and gauge bosons [J]. Physics, 2015; arXiv:1507.02330.
- [22] DAVOUDIASHVILI H, LEE H S, MARCIANO W J. Dark side of higgs diphoton decays and muon $g - 2$ [J]. Physical Review D, 2012, 86(9): 095009(1-7).
- [23] ENDO M, HAMAGUCHI K, MISHIMA G. Constraints on hidden photon models from electron $g - 2$ and hydrogen spectroscopy [J]. Physical Review D, 2012, 86 (9): 579-599.
- [24] GNINENKO S N, KRASNIKOV N V. The muon anomalous magnetic moment and a new light gauge boson [J]. Physics Letters B, 2001, 513(1-2): 119-122.
- [25] FAYET P. U -boson production in $e^+ e^-$ annihilations, ψ and γ decays, and light dark matter [J]. Physical Review D, 2007, 75(11): 115017(1-18).
- [26] POSPELOV M. Secluded $U(1)$ below the weak scale [J]. Physical Review D, 2009, 80(9): 281-287.
- [27] ABRAHAMYAN S, AHMED Z, ALLADA K, et al. Search for a new gauge boson in electron-nucleus fixed-target scattering by the APEX experiment [J]. Physical Review Letters, 2011, 107(19): 191804(1-5).
- [28] MERKEL H, ACHENBACH P, GAYOSO C A, et al. Search for light massive gauge bosons as an explanation of the $(g - 2)_\mu$ anomaly at MAMI [J]. Physical Review Letters, 2014, 112: 221802(arXiv:1404.5502).
- [29] ANDREASS, NIEBUHR C, RINGWALD A. New limits on hidden photons from past electron beam dumps [J]. Physical Review D, 2012, 86(9): 095019(arXiv:1209.6083).
- [30] BJORKEN J D, ESSIG R, SCHUSTER P, et al. New fixed-target experiments to search for dark gauge forces [J]. Physical Review D, 2009, 80(7): 264-284.
- [31] BABUSCI D, BADONI D, BALWIERZ-PYTKO I, et al. Limit on the production of a light vector gauge boson in phi meson decays with the KLOE detector [J]. Physics Letters B, 2013, 720(1-3): 111-115.
- [32] AGAKISHIEV G, BALANDA A, BELYAEV A, et al. Searching a dark photon with HADES [J]. Physics Letters B, 2014, 731(2): 265-271(arXiv:1311.0216).
- [33] ADARE A, AFANASIEV S, AIDALA C, et al. Search for dark photons from neutral meson decays in $p^+ p$ and $d^+ Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV [J]. Physical Review C, 2015, 91(3): 031901(arXiv:1409.0851).
- [34] GUNJI T. Dark Photon Searches at ALICE [R]. Presented at Dark Interactions 2014 Workshop at Brookhaven National Laboratory, USA.
- [35] BATLEY J R, KALMUS G, LAZZERONI C, et al. Search for the dark photon in π^0 decays [J]. Physics Letters B, 2015, 746: 178-185(arXiv:1504.00607).
- [36] BABUSCI D, BALWIERZ-PYTKO I, BENCIVENNI G, et al. Search for light vector boson production in $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$ interactions with the KLOE experiment [J]. Physics Letters B, 2014, 736: 459(arXiv:1404.7772).
- [37] LEES J P, POIREAU V, TISSERAND V, et al. Search for a dark photon in $e^+ e^-$ collisions at BABAR [J]. Physical Review Letters, 2014, 113(20): 201801 (arXiv:1406.2980).
- [38] ERLER J, RAMSEY-MUSOLF M J. Weak mixing angle at low energies [J]. Physical Review D, 2005, 72: 073003 (arXiv:hep-ph/0409169).
- [39] JEGERLEHNER F. Electroweak effective couplings for future precision experiments [J]. Eprint Arxiv, 2011, 31; arXiv:1107.4683.
- [40] BURGER F, JANSEN K, PETSCHLIES M, et al. Leading hadronic contributions to the running of the electroweak coupling constants from lattice QCD [J]. Journal of High Energy Physics, 2015, 11 (11): 1-23 (arXiv:1505.03283).
- [41] FRANCIS A, GÜLPERS V, HERDÓZA G, et al. Study of the hadronic contributions to the running of the QED coupling and the weak mixing angle [J]. Physics, 2015,

- 110: 101-103(arXiv:1511.04751).
- [42] GILBERT S L, NOECKER M C, WATTS R N, et al. Measurement of parity nonconservation in atomic cesium [J]. *Physical Review Letters*, 1985, 55(24): 2680 (doi.org/10.1103/PhysRevLett.55.2680).
- [43] WOOD C S, BENNETT S C, CHO D, et al. Measurement of parity nonconservation and an anapole moment in cesium[J]. *Science*, 1997, 275(5307): 1759-1763.
- [44] BENNETTS C, WIEMAN C E. Erratum: Measurement of the $6s \rightarrow 7s$ transition polarizability in atomic cesium and an improved test of the standard model [J]. *Physical Review Letters*, 1999, 82(12): 2484-2487.
- [45] PORTELA M N, DIJCK E A, MOHANTY A, et al. Ra^+ ion trapping: Toward an atomic parity violation measurement and an optical clock [J]. *Applied Physics B*, 2014, 114: 173-182.
- [46] JUNGSMANN K P. Symmetries and fundamental interactions-selected topics [J]. *Hyperfine Interactions*, 2014, 227(1): 5-16.
- [47] ANTHONY P L, ARNOLD R G, ARROYO C, et al. Precision measurement of the weak mixing angle in Moller scattering[J]. *Physical Review Letters*, 2005, 95(8): 081601(hep-ex/0504049).
- [48] ANDROIC D, ARMSTRONG D S, ASATURYAN A, et al. First determination of the weak charge of the proton [J]. *Physical Review Letters*, 2013, 111(14): 141803 (arXiv:1307.5275).
- [49] MAMMEI J. The MOLLER Experiment [J]. *Physics*, 2012, 35(4): 203(arXiv:1208.1260).
- [50] AULENBACHER K. Opportunities for parity violating electron scattering experiments at the planned MESA facility[J]. *Hyperfine Interactions*, 2011, 200(1): 3-7.
- [51] WANG D, PAN K, SUBEDI R, et al. Measurement of parity violation in electron-quark scattering [J]. *Nature*, 2014, 506(7486): 67-70.
- [52] REIMER P E. Parity violating deep inelastic scattering [J]. *Nuovo Cimento C*, 2012, 35(4): 209-213.
- [53] ZELLER G P, Mcfarland K S, ADAMS T, et al. Precise determination of electroweak parameters in neutrino-nucleon scattering [J]. *Physical Review Letters*, 2002, 88(9): 386-389.